



Responding to Suicidal Ideation

Suicidal Ideation:	Thinking about committing suicide. Passive suicidal ideation is when you wish were dead or that you could die, but you don't actually have any plan. Active suicidal ideation, on the other hand, is not only thinking about it but having the intent to commit suicide, including planning how to do it.
---------------------------	---

Overview of Responding to Suicidal Ideation:

Clients identifying that they're struggling with suicidal thoughts is one of the more intimidating and challenging situations for a residential counselor. In general, the empathic listening technique can be very useful for discussing suicidal ideation with the client.

By way of review Empathic Listening involves:

- Mirror or match the client's general body language, speech, intensity; lead to slower and lower.
- Intense Listening.
- Asking clarifying questions in a non-judgmental fashion.
- Focusing on emotions.
- Avoid slipping into problem-solving.
- Avoid the "empathizing equals agreeing with" trap if client is being negative about others.
- Plug the client back into program structures.

The intent of Empathic Listening is for the client to experience a feeling of being understood and supported by staff. It is not a problem-solving technique. Rather than attempting to fix *why* the client is having suicidal thoughts, instead focus on helping them see those thoughts as a symptom. How often does this happen? How strong are these thoughts? How intrusive are they? When are they happening? What triggers having these sorts of thoughts?

The Columbia Protocol suggests a series of questions to determine the extent, if any, to which the client has a plan or has taken any actions toward suicide.

1. Have you wished you were dead or wished you could go to sleep and not wake up?
2. Have you been thinking about killing yourself?
3. Have you been thinking about how you might do this?
4. Have you had these thoughts and had some intention of acting on them?
5. Have you started to work out or worked out the details of how to kill yourself? Did you intend to carry out this plan?
6. Have you done anything, started to do anything, or prepared to do anything to end your life (such as collecting pills, giving valuables away, writing a suicide note, etc.)?

Ask the client if they have any contraband items to turn in to staff.



Suicidal Gesture:	A non-verbal behavior expressing suicidal intent.
--------------------------	---

A common way to assess suicidal gestures or suicide plans is to evaluate both the risk factors inherent in the attempt and the rescue factors.

Risk / Rescue Assessment for suicidal gestures:

A. Method of attempt:

1. No or questionable seriousness: nonlethal, recovery certain.
2. Mildly serious: Death is unlikely, but possible as a secondary complication.
3. Moderately serious: Possibility of death.
4. Highly serious: Death is likely.
5. Extremely serious: Death is the usual outcome; survival chance is minimal to none.

B. Rescuability:

1. High: Most easily rescued.
2. High / Moderate: More chance of rescue.
3. Moderate: Chance of rescue 50:50.
4. Low / Moderate: Less chance of rescue.
5. Low: Chance of rescue is minimal to none.

C. Physical Consequences:

1. No or nonspecific symptoms.
2. Mild symptoms.
3. Moderate symptoms.
4. Severe symptoms.
5. Extremely severe symptoms, organ failures.

E. Medical intervention need:

1. Medical intervention not required or observation only.
2. First aid.
3. Emergency care.
4. In-patient care, specialized units.
5. Intensive-care, life support.

F. Global impression of lethality:

1. Subliminal: Death impossible or highly improbable.
2. Low: Death improbable.
3. Moderate: Death probable.
4. High: Death highly probable.
5. Extremely high: Death extremely probable to almost certain.



Bear Clan, LLC

Empowering Balanced and Transformative Lives

The following chart presents a slightly different way to look at risk factors:

Talking About:	Behaviors	Mood
<ul style="list-style-type: none">• Killing themselves;• Feeling hopeless;• Having no reason to live;• Being a burden to others;• Feeling trapped;• Unbearable pain.	<ul style="list-style-type: none">• Increased use of alcohol or drugs;• Looking for a way to end their lives, such as searching online for methods;• Withdrawing from activities;• Isolating from family and friends;• Sleeping too much or too little;• Visiting or calling people to say goodbye;• Giving away prized possessions;• Idolizing celebrities who have committed suicide;• Aggression;• Fatigue.	<ul style="list-style-type: none">• Depression;• Anxiety;• Loss of interest;• Irritability;• Humiliation/Shame;• Agitation/Anger;• Relief/Sudden Improvement.

Rescue factors frequently involve the person letting other people know about their suicidal ideation or gesture, or making comments or gestures in a location where their distress is going to be noticed by responsible adults.

Suicidal gestures can be related to self-injurious behaviors (such as superficial cutting or head-banging). However, self-injurious behaviors typically are not directly correlated with suicidality; are most often seen as examples of non-suicidal self-injury, serving a different psychological purpose. Nevertheless, self-injurious behaviors should be reported to program leadership, a clinician, or whatever protocols your program uses, as soon as possible.

Qualities to look for in accessing risk factors include:

- **Excessive sadness or moodiness:** Long-lasting sadness, mood swings, and unexpected rage.
- **Hopelessness:** Feeling a deep sense of hopelessness about the future, with little expectation that circumstances can improve.
- **Sleep problems.**
- **Sudden calmness:** Suddenly becoming calm after a period of depression or moodiness can be a sign that the person has made a decision to end his or her life.
- **Withdrawal:** Choosing to be alone and avoiding friends or social activities also are possible symptoms of depression, a leading factor in suicide. This includes the loss of interest or pleasure in activities the person previously enjoyed.
- **Changes in personality and/or appearance:** A person who is considering suicide might exhibit a change in attitude or behavior, such as speaking or moving with unusual speed or



Bear Clan, LLC

Empowering Balanced and Transformative Lives

slowness. In addition, the person might suddenly become less concerned about his or her personal appearance.

- **Dangerous or self-harmful behavior:** Potentially dangerous behavior, such as reckless driving, engaging in unsafe sex, and increased use of drugs and/or alcohol might indicate that the person no longer values his or her life.
- **Recent trauma or life crisis:** A major life crises might trigger a suicide attempt. Crises include the death of a loved one or pet, divorce or break-up of a relationship, diagnosis of a major illness, loss of a job, or serious financial problems.
- **Making preparations:** Often, people considering suicide will begin to put their personal business in order. This might include visiting friends and family members, giving away personal possessions, making a will, and cleaning up their room or home. Some people will write a note before committing suicide. Some will buy a firearm or other means like poison.
- **Threatening suicide:** 50% to 75% of those considering suicide will give someone – a friend or relative – a warning sign. However, not everyone who is considering suicide will say so, and not everyone who threatens suicide will follow through with it. Every threat of suicide must be taken seriously.

After processing suicidal ideation:

Communication and Initial Documentation (will be different in different programs):

1. Contact Health Services if there is any medical follow up indicated.
2. Verbally communicate about the incident of suicidal ideation to your shift colleagues and to staff covering the next shift.
3. Contact the On-Call Clinician.
4. Document the incident on an Incident Report Form. Your processing of the suicidal ideation may not be captured fully by the Incident Report so if indicated send a separate email to the resident's clinician and/or case manager with additional information.
5. Complete other documentation used by your program such as a Daily Shift Log.

Structural Interventions:

In general, residents should be returned to the regular program structures. However, similar to after any emotionally intense incident, it may make sense to add some additional structures.

Possibilities include, but are not limited to:

- ☒ Scheduled Check-In's for the remainder of the shift
- ☒ Increasing structure of various activities – for example, by only giving one-step directions.
- ☒ Staff Shadowing – requiring the resident to remain in arms reach of staff.
- ☒ Re-entry Plan – requiring a meeting to facilitate the resident being reintegrated with the group.
- ☒ Relationship Repair activities and/or Restitution
- ☒ Safety Watch (see Safety Watch form).
- ☒ Safety Contract (see below).